




NF- κ B p65 (Phospho-Ser468) Antibody

E011013

- Catalog Number:** E011013-1, E011013-2
Amount: 50 μ g/50 μ l, 100 μ g/100 μ l
Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage/Stability: Store at -20°C/1 year
Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human NF- κ B around the phosphorylation site of serine 468 (L-A-S^P-V-D).
Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.
Specificity/Sensitivity: NF- κ B p65 (phospho-Ser468) antibody detects endogenous levels of NF- κ B p65 only when phosphorylated at serine 468.
Reactivity: Human, Mouse, Rat
Applications: WB: 1:500~1:1000 IF:1:100~1:200
Swiss-Prot No. : Q04206
References: Buss H, et al. (2004) J Biol Chem.
Schmitz ML, et al. (2004) Blood.



For Research Use Only